



Deductive Reasoning Based on the Valid Aristotelian Syllogism *AEE-2*



Haiping Wang*

School of Philosophy, Anhui University, China

Email: 1106764362@qq.com



Xiaojun Zhang

School of Philosophy, Anhui University, China

Email: 591551032@qq.com

Article History

Received: 4 January, 2024

Revised: 25 March, 2024

Accepted: 27 March, 2024

Published: 29 March, 2024

How to Cite

Haiping Wang, Xiaojun Zhang (2024). Deductive Reasoning Based on the Valid Aristotelian Syllogism *AEE-2*. *Sumerianz Journal of Social Science*, 7(1): 1-4.

Abstract

This paper firstly formalizes Aristotelian syllogisms based on the tripartite structure of categorical propositions, and then uses the truth definitions of categorical propositions to prove the validity of the Aristotelian syllogism *AEE-2*. Then, the remaining 23 valid syllogisms are derived from the syllogism *AEE-2* with the help of relevant facts, inner and outer negation definitions of quantifiers, and deductive rules. In other words, this paper reveals the reducible relationship between/among these 24 syllogisms and establishes a succinct formal reason system for Aristotelian syllogistic. The deductive reasoning not only ensures consistency in its results, but also provides a concise mathematical paradigm for other types of syllogisms.

Keywords: Aristotelian syllogisms; Deductive reasoning; Reducible relationship; Validity.

1. Introduction

In natural language, there are many types of syllogisms, such as Aristotelian syllogisms (Yijiang, 2023), modal syllogisms (Cheng, 2023), generalized syllogisms (Moss, 2010), and so on, which are common and important forms of reasoning in social life and logic (Jing and Xiaojun, 2023), and has been widely studied since Aristotle (Murinová and Novák, 2012). This paper focuses on the study of Aristotelian syllogisms. And in the following, unless otherwise specified, syllogisms refer to Aristotelian syllogisms.

Aristotelian syllogisms involve the following four types of statements: *all rs are t*, *no rs are t*, *some rs are t*, and *not all rs are t*. They are abbreviated as Propositions *A*, *E*, *I*, and *O* respectively, where *all*, *no*, *some*, and *not all* are called Aristotelian quantifiers (Long and Xiaojun, 2023). It is known that there are only 24 Aristotelian syllogisms out of 256 ones (Xiaojun *et al.*, 2022).

Lukasiewicz (1957), took the syllogisms *AAA-1* and *AII-3* as basic axioms to derive the other 22 valid ones. Xiaojun and Sheng (2016), deduced the other 22 valid syllogisms on the basis of the two syllogisms *AAA-1* and *EAE-1*. This paper only uses the syllogism *AEE-2* as a basic axiom reasoning basis to infer the remaining 23 valid ones.

2. Formal Aristotelian Syllogistic

In order to construct the formal system, the following four parts need to be provided: (1) primitive symbols; (2) formation rules of well-formed formulas (abbreviated as wff); (3) basic axioms; (4) rules of inference. In the following, let α , β , γ and δ be wffs.

2.1. Primitive Symbols

- lexical variables: r , s , t
- negative operator: \neg

- implication operator: \rightarrow
- quantifier: *all*
- brackets: (,)

2.2. Formation Rules

- If Q is a quantifier, r and t are lexical variables, then $Q(r, t)$ is a wff.
- If α and β are wffs, then $\neg\alpha$ and $\alpha\rightarrow\beta$ are wffs.
- Only the formulas obtained based on (2.2.1) and (2.2.2) are wffs.

For instance, $all(r, t)$, and $all(r, t)\rightarrow some(r, s)$ are wffs that can be seen as ‘all rs are t ’ and ‘if all rs are t , then some rs are s ’, respectively. The others are similar. It can be known that Aristotelian syllogisms contain the following four types of categorical propositions, namely $all(r, t)$, $no(r, t)$, $some(r, t)$, and $not\ all(r, t)$. They are respectively abbreviated as Propositions A, E, I and O .

2.3. Basic Axioms

A1: If α is a valid formula in first-order logic, then $\vdash\alpha$.

A2: $\vdash all(t, s)\wedge no(r, s)\rightarrow no(r, t)$ (that is, the syllogism $AEE-2$).

In these axioms, ‘ $\vdash\alpha$ ’ means that α is provable. The other cases are similar.

2.4. Rules of Inference

R1: (subsequent weakening): From $\vdash(\alpha\wedge\beta\rightarrow\gamma)$ and $\vdash(\gamma\rightarrow\delta)$, infer $\vdash(\alpha\wedge\beta\rightarrow\delta)$.

R2: (anti-syllogism): From $\vdash(\alpha\wedge\beta\rightarrow\gamma)$, infer $\vdash(\neg\gamma\wedge\alpha\rightarrow\neg\beta)$.

2.5. Related Definitions

D1 (biconditional \leftrightarrow): $(\alpha\Box\beta)=_{\text{def}}(\alpha\rightarrow\beta)\wedge(\beta\rightarrow\alpha)$.

D2: $(\alpha\wedge\beta)=_{\text{def}}\neg(\alpha\rightarrow\neg\beta)$.

D3 (inner negative quantifier): $Q\neg(r, t)=_{\text{def}}Q(r, D\neg t)$.

D4 (outer negative quantifier): $(\neg Q)(r, t)=_{\text{def}}$ It is not that $Q(r, t)$.

D5 (true value of quantifier): $all(r, t)=_{\text{def}}R\subseteq T$.

D6 (true value of quantifier): $some(r, t)=_{\text{def}}R\cap T\neq\emptyset$.

D7 (true value of quantifier): $no(r, t)=_{\text{def}}R\cap T=\emptyset$.

D8 (true value of quantifier): $not\ all(r, t)=_{\text{def}}R\nsubseteq T$.

2.6. Related Facts

Fact 1 (inner negation):

F1: $\vdash all(r, t)\leftrightarrow no\neg(r, t)$; F2: $\vdash no(r, t)\leftrightarrow all\neg(r, t)$;

F3: $\vdash some(r, t)\leftrightarrow not\ all\neg(r, t)$; F4: $\vdash not\ all(r, t)\leftrightarrow some\neg(r, t)$.

Fact 2 (outer negation):

F5: $\vdash all(r, t)\Box not\ all(r, t)$; F6: $\vdash not\ all(r, t)\Box all(r, t)$;

F7: $\vdash some(r, t)\Box no(r, t)$; F8: $\vdash no(r, t)\Box some(r, t)$.

Fact 3 (symmetry of *some* and *no*):

F9: $\vdash some(r, t)\Box some(t, r)$; F10: $\vdash no(r, t)\Box no(t, r)$.

Fact 4 (assertoric subalternations):

F11: $\vdash no(r, t)\Box not\ all(r, t)$; F12: $\vdash all(r, t)\Box some(r, t)$.

3. Reducible Relationship between the Other 23 Valid Syllogisms and the Syllogism $AEE-2$

The following Theorem 1 shows the syllogism $AEE-2$ is valid. ‘ $AEE-2\Box\Box\Box\Box\Box\Box$ in Theorem 2 indicates that the validity of the syllogism $\Box\Box\Box\Box\Box\Box$ can be derived from the validity of the syllogism $AEE-2$. That is to say, there is a reducible relationship between these two syllogisms. The others are similar.

Theorem 1 ($AEE-2$): $\vdash all(t, s)\wedge no(r, s)\rightarrow no(r, t)$ is valid.

Proof: Suppose that $all(t, s)$ and $no(r, s)$ are true, then $T\subseteq S$ and $R\cap S=\emptyset$ are true according to Definition 5 and 7, respectively. Hence, it can be seen that $R\cap T=\emptyset$ is true. Thus $no(r, t)$ is true in line with Definition 7. It follows that $\vdash all(t, s)\wedge no(r, s)\rightarrow no(r, t)$ is valid, as required.

Theorem 2: The remaining 23 valid syllogisms can be deduced just from the syllogism $AEE-2$:

(1) $\vdash AEE-2\Box AEE-4$

(2) $\vdash AEE-2\Box AEE-4\Box EAE-1$

(3) $\vdash AEE-2\Box EAE-2$

(4) $\vdash AEE-2\Box AII-1$

(5) $\vdash AEE-2\Box AII-1\Box AII-3$

(6) $\vdash AEE-2\Box AII-1\Box AII-3\Box IAI-3$

(7) $\vdash AEE-2\Box AII-1\Box IAI-4$

(8) $\vdash AEE-2\Box AEO-2$

- (9) $\vdash AEE-2 \square AEO-2 \square AEO-4$
- (10) $\vdash AEE-2 \square AEE-4 \square EAE-1 \square EAO-1$
- (11) $\vdash AEE-2 \square AEE-4 \square EAE-1 \square EAO-1 \square EAO-2$
- (12) $\vdash AEE-2 \square AEE-4 \square EAE-1 \square EAO-1 \square EAO-2 \square AAI-3$
- (13) $\vdash AEE-2 \square AEO-2 \square AAI-1$
- (14) $\vdash AEE-2 \square AEO-2 \square AAI-1 \square AAI-4$
- (15) $\vdash AEE-2 \square AEO-2 \square AAI-1 \square AAI-4 \square EAO-4$
- (16) $\vdash AEE-2 \square AEO-2 \square AAI-1 \square AAI-4 \square EAO-4 \square EAO-3$
- (17) $\vdash AEE-2 \square AEE-4 \square EAE-1 \square AAA-1$
- (18) $\vdash AEE-2 \square AEE-4 \square EAE-1 \square AAA-1 \square OAO-3$
- (19) $\vdash AEE-2 \square AEE-4 \square EAE-1 \square AAA-1 \square OAO-3 \square AOO-2$
- (20) $\vdash AEE-2 \square AII-1 \square EIO-1$
- (21) $\vdash AEE-2 \square AII-1 \square EIO-1 \square EIO-3$
- (22) $\vdash AEE-2 \square AII-1 \square EIO-1 \square EIO-3 \square EIO-4$
- (23) $\vdash AEE-2 \square AII-1 \square EIO-1 \square EIO-2$

Proof:

- | | |
|--|---|
| [1] $\vdash all(t, s) \wedge no(r, s) \square no(r, t)$ | (i.e. <i>AEE-2</i> , basic axiom A2) |
| [2] $\vdash all(t, s) \wedge no(s, r) \square no(r, t)$ | (i.e. <i>AEE-4</i> , by [1] and F10) |
| [3] $\vdash all(t, s) \wedge no(s, r) \square no(t, r)$ | (i.e. <i>EAE-1</i> , by [2] and F10) |
| [4] $\vdash all(t, s) \wedge no(r, s) \square no(t, r)$ | (i.e. <i>EAE-2</i> , by [1] and F10) |
| [5] $\vdash \neg no(r, t) \wedge all(t, s) \square \neg no(r, s)$ | (by [1] and R2) |
| [6] $\vdash some(r, t) \wedge all(t, s) \square some(r, s)$ | (i.e. <i>AII-1</i> , by [5] and F7) |
| [7] $\vdash some(t, r) \wedge all(t, s) \square some(r, s)$ | (i.e. <i>AII-3</i> , by [6] and F9) |
| [8] $\vdash some(t, r) \wedge all(t, s) \square some(s, r)$ | (i.e. <i>IAI-3</i> , by [7] and F9) |
| [9] $\vdash some(r, t) \wedge all(t, s) \square some(s, r)$ | (i.e. <i>IAI-4</i> , by [6] and F9) |
| [10] $\vdash no(r, t) \square not\ all(r, t)$ | (by F11) |
| [11] $\vdash all(t, s) \wedge no(r, s) \square not\ all(r, t)$ | (i.e. <i>AEO-2</i> , by [1], [10] and R1) |
| [12] $\vdash all(t, s) \wedge no(s, r) \square not\ all(r, t)$ | (i.e. <i>AEO-4</i> , by [11] and F10) |
| [13] $\vdash all(t, s) \wedge no(s, r) \square not\ all(t, r)$ | (i.e. <i>EAO-1</i> , by [3], [10] and R1) |
| [14] $\vdash all(t, s) \wedge no(r, s) \square not\ all(t, r)$ | (i.e. <i>EAO-2</i> , by [13] and F10) |
| [15] $\vdash \neg not\ all(t, r) \wedge all(t, s) \square \neg no(r, s)$ | (by [14] and R2) |
| [16] $\vdash all(t, r) \wedge all(t, s) \square some(r, s)$ | (i.e. <i>AAI-3</i> , by [15], F5 and F7) |
| [17] $\vdash \neg not\ all(r, t) \wedge all(t, s) \square \neg no(r, s)$ | (by [11] and R2) |
| [18] $\vdash all(r, t) \wedge all(t, s) \square some(r, s)$ | (i.e. <i>AAI-1</i> , by [17], F5 and F7) |
| [19] $\vdash all(r, t) \wedge all(t, s) \square some(s, r)$ | (i.e. <i>AAI-4</i> , by [18] and F9) |
| [20] $\vdash \neg some(s, r) \wedge all(r, t) \square \neg all(t, s)$ | (by [19] and R2) |
| [21] $\vdash no(s, r) \wedge all(r, t) \square not\ all(t, s)$ | (i.e. <i>EAO-4</i> , by [20], F8 and F6) |
| [22] $\vdash no(r, s) \wedge all(r, t) \square not\ all(t, s)$ | (i.e. <i>EAO-3</i> , by [21] and F10) |
| [23] $\vdash all(t, s) \wedge all \neg(s, r) \square all \neg(t, r)$ | (by [3] and F2) |
| [24] $\vdash all(t, s) \wedge all(s, D \neg r) \square all(t, D \neg r)$ | (i.e. <i>AAA-1</i> , by [23] and D3) |
| [25] $\vdash \neg all(t, D \neg r) \wedge all(t, s) \square \neg all(s, D \neg r)$ | (by [24] and R2) |
| [26] $\vdash not\ all(t, D \neg r) \wedge all(t, s) \square not\ all(s, D \neg r)$ | (i.e. <i>OAO-3</i> , by [25] and F6) |
| [27] $\vdash \neg not\ all(s, D \neg r) \wedge not\ all(t, D \neg r) \square \neg all(t, s)$ | (by [26] and R2) |
| [28] $\vdash all(s, D \neg r) \wedge not\ all(t, D \neg r) \square not\ all(t, s)$ | (i.e. <i>AOO-2</i> , by [27], F5 and F6) |
| [29] $\vdash some(r, t) \wedge no \neg(t, s) \square not\ all \neg(r, s)$ | (by [6], F1 and F3) |
| [30] $\vdash some(r, t) \wedge no(t, D \neg s) \square not\ all(r, D \neg s)$ | (i.e. <i>EIO-1</i> , by [29] and D3) |
| [31] $\vdash some(t, r) \wedge no(t, D \neg s) \square not\ all(r, D \neg s)$ | (i.e. <i>EIO-3</i> , by [30] and F9) |
| [32] $\vdash some(t, r) \wedge no(D \neg s, t) \square not\ all(r, D \neg s)$ | (i.e. <i>EIO-4</i> , by [31] and F10) |
| [33] $\vdash some(r, t) \wedge no(D \neg s, t) \square not\ all(r, D \neg s)$ | (i.e. <i>EIO-2</i> , by [30] and F10) |

So far, on the basis of 33 reasoning steps, Theorem 2 has completed the task of transforming the syllogism *AEE-2* to the other 23 valid syllogisms.

4. Conclusion

This paper firstly formalizes Aristotelian syllogisms based on the tripartite structure of categorical propositions, and then uses the truth definitions of categorical propositions to prove the validity of the Aristotelian syllogism *AEE-2*. Then, the remaining 23 valid syllogisms are derived from the syllogism *AEE-2* with the help of relevant facts, inner and outer negation definitions of quantifiers, and deductive rules. In other words, this paper reveals the reducible relationship between/among these 24 syllogisms and establishes a succinct formal reason system for Aristotelian syllogistic.

The deductive reasoning not only ensures consistency in its results, but also provides a concise mathematical paradigm for other types of syllogisms (such as generalized modal syllogisms, syllogisms with adjectives). How to implement this formal method on a computer? This question requires further study.

Acknowledgements

This work was supported by the National Social Science Foundation of China under Grant No.21BZX100.

References

- Cheng, Z. (2023). How to deduce the other 91 valid aristotelian modal syllogisms from the syllogism iai-3. *Applied Science and Innovative Research*, 7(1): 46-57.
- Jing, X. and Xiaojun, Z. (2023). The reducibility of generalized modal syllogisms based on ami-1. *SCIREA Journal of Philosophy*, 3(1): 1-11.
- Long, W. and Xiaojun, Z. (2023). The reducibility of modal syllogisms based on the syllogism ei \diamond o-2. *SCIREA Journal of Mathematics*, 8(3): 87-96.
- Łukasiewicz, J. (1957). *Aristotle's syllogistic: From the standpoint of modern formal logic*. Second edn: Clarendon Press: Oxford.
- Moss, L. S. (2010). Syllogistic logics with verbs. *Journal of Logic and Computation*, 20(4): 947-67.
- Murinová, P. and Novák, V. (2012). A formal theory of generalized intermediate syllogisms. *Fuzzy Sets and Systems*, 186: 47-80. Available: <https://doi.org/10.1016/j.fss.2011.07.004>.
- Xiaojun, Z. and Sheng, L. (2016). Research on the formalization and axiomatization of traditional syllogisms. *Journal of Hubei University (Philosophy and Social Sciences)*, 6: 32-37. Available: <https://www.scirp.org/reference/referencespapers?referenceid=3454840>
- Xiaojun, Z., Hui, L. and Yijiang, H. (2022). How to deduce the remaining 23 valid syllogisms from the validity of the syllogism eio-1. *Applied and Computational Mathematics*, 11(6): 160-64.
- Yijiang, H. (2023). The reductions between/among aristotelian syllogisms based on the syllogism aii-3. *SCIREA Journal of Philosophy*, 3(1): 12-22.